arXiv:math/0102068v1 [math.NT] 8 Feb 2001

RAMIFICATION IN C-EXTENSIONS OF LOCAL FIELDS OF CHARACTERISTIC 0

Odile Sauzet

Abstract. We construct explicitly APF extensions of finite extensions of $\mathbb{Q}_p$ for which the Galois group is not a p-adic Lie group and which do not have any open subgroup with $\mathbb{Z}_p$-quotient.

Let K be a finite extension of $\mathbb{Q}_p$ and L/K be a Galois totally ramified pro-p-extension. If the Galois group of L/K is a p-adic Lie group it is known from Sen ([Sen]) that the sequence of upper ramification breaks of L/K is unbounded and that the higher ramification groups are open in $Gal(L/K)$, i.e. in the terminology of [FW], the extension is arithmetically profinite (APF).

The more general question concerning the existence of APF extensions of finite extensions of $\mathbb{Q}_p$ with a Galois group which is a free pro-p-group remains open (the answer is positive in characteristic p, see [Fe]) and we are interested in obtaining methods to construct APF extensions with a non p-adic Lie Galois group.

Coates and Greenberg ([CG]) raised the question whether the notion of unbounded sequence of upper ramification breaks (extensions deeply ramified) depends on the existence of an open subgroup of the Galois group having a quotient isomorphic to $\mathbb{Z}_p$. Fesenko answered this question negatively in [Fe] by showing the existence of a finitely generated pro-p-group T such that no open subgroup contains a $\mathbb{Z}_p$-quotient. He showed that the group T can be realized a the Galois group of a finite extension of $\mathbb{Q}_p$.

In this paper we look at pro-p-groups such that the lower p-series define a filtration of open subgroups. We show that this type of group can be realized as the Galois group of a deeply ramified extension (and arithmetically profinite). Not only the reason for being deeply ramified is not due to the presence of a $\mathbb{Z}_p$-extension but we show that this notion is independent of the nature of the Galois group, i.e. there are totally ramified extensions with such a Galois group which are not deeply ramified.

We give a method to construct explicitly APF extensions with a Galois group such as we described above. The method consists in taking a tower of extension K_i/K_{i-1} such that K_{i+2}/K_{i-1} is a Galois extension of order p^3 with Galois group having two generators a_i and a_{i+1} and the Galois group of K_{i+2}/K_{i+1} is generated by the commutator $[a_{i+1}, a_i]$. We show how we can control asymptotically the ramification in $M = \cup_i K_i/K$. We call the normal closure of such an extension a C-extension (the "C" standing for commutators)

Supported by a Marie Curie Fellowship from the European Union

After recalling some background on ramification in Section 0, we give results on the structure of the Galois group of the normal closure of the extension L/K which are relevant to the identification of the sequence of ramification breaks. Namely, we prove that the Galois group does not contain any proper closed normal subgroup. Therefore a C-extension is deeply ramified if and only if it is APF. And we show that the group does not contain any open subgroup with a $\mathbb{Z}_p$-quotient.

In Section 2 we construct the tower described above and we show how we can have it to be APF. In Section 3 we show that the unboundedness of the ramification breaks does not depend on the structure of the Galois group because there are C-extensions which are not deeply ramified. The last section shows how one can use C-extensions to construct more APF extensions. Working on problems relative to arithmetically profinite extensions was suggested to me by Ivan Fesenko. I am grateful for his support and numerous comments.

0. Ramification and arithmetically profinite extensions.

We recall some basic notion of ramification theory, for finite extensions see [Ser], and for infinite extensions see [FV]. For a general introduction to APF extensions see [DuFe].

Let L/K be a totally ramified finite Galois p-extension of local fields with finite residue field and with Galois group G. Let

$$i_G(\sigma) = v_L(\frac{\sigma\pi_L}{\pi_L} - 1);$$

where π_L is a uniformiser of L and σ an element in G.

The ramification filtration in the lower numbering of G is defined by (see [Ser1, Chap. VI])

$$G_i = \{\sigma \in G \ ; \ i_G(\sigma) \geq i+1\}.$$

This filtration does not behave well with quotients. To define ramification for infinite extensions, there is another filtration defined via the Herbrand function $h_{L/K}$ whose explicit formulation in the case of totally ramified p-extensions is as follow :

$$h_{L/K}(x) = \begin{cases} x & \text{if } \ x \leq i \\ (p-1)i + px & \text{otherwise} \end{cases} \quad \text{if } [L:K] = p \ ;$$

and for a general finite extension the function is obtained with the equality :

$$h_{L/K} = h_{L/M} \circ h_{M/K}$$

where M/K is a subextension of L/K.

Let consider a group G of wild automorphisms of a local field of characteristic p with finite residue field, we can define the ramification filtration in a similar way, we write h_G for the Herbrand function.

Then the ramification filtration in the upper numbering is given by

$$G^u = G$$

Let H be a normal subgroup of G, then

$$(G/H)^u = G^u.H/H.$$

Now we can define the upper ramification filtration for infinite p-extensions :

Let L/K be an infinite Galois extension such that $L = \cup_i K_i$, then

$$G = Gal(L/K) = \varprojlim Gal(K_i/K)$$

and

$$G^u = \varprojlim Gal(K_i/K)^u.$$

Definition 0.1. *The totally ramified extension L/K is arithmetically profinite (APF) if for all $x \geq 0$, the normal subgroup G^x is non trivial and open in G*

If the extension L/K is APF, then we can again define the Herbrand function :

$$h_{L/K}(x) = \lim_{i \to \infty} h_{K_i/K}(x),$$

and the ramification filtration of G in the lower numbering is well defined.

Fontaine and Wintenberger have defined the field of norms $N(L/K)$ for an APF extension L/K ([FW]) :

$$N(L/K)^\times = \varprojlim K_i^\times \quad ; \quad N(L/K) = N(L/K)^\times \cup \{0\}$$

where $\{K_i\}$ is an ordered family of finite subextensions of L/K such that $L = \cup K_i$ and the connecting maps are the norm. If L/K is APF, then the valuation defined in [W2, 2.1.2.], addition defined in [W2, 2.1.3.] and the natural multiplication give to $N(L/K)$ a structure of local field of characteristic p with residue field isomorphic to the the one of K and we have the following result on the action of G on $N(L/K)$ and the ramification ([W2, 3.3.4.]) :

Proposition 0.2. *G acts faithfully on $N(L/K)$ and can be identified to a closed subgroup of $Aut(N(L/K))$. Moreover the ramification filtration of G can be read in its image in $Aut(N(L/K))$, i.e. if $N_{L/K}(\sigma)$ is the image of an element $\sigma \in G$ in $Aut(N(L/K))$, then*

$$G_i = \{\sigma \in G \ ; \ i_{N(L/K)}(N_{L/K}(\sigma)) \geq i\}.$$

The following notations will be used in the sequel :

- The ramification break i of a finite p-extension or an APF p-extension with Galois group G, or a group of automorphism G of a local field of characteristic p is defined by

$$i = \sup\{x > 0, G = G_x\}.$$

- The upper ramification break of G is $u(G) = h_{L/K}^{-1}(i(G))$.

1. Definitions and basic properties

A pro-p-extension L/K can be defined as a tower of extensions K_n/K such that $[K_n : K_{n-1}] = p$. If the extension L/K is Galois they correspond to a central series of the Galois group. In order to construct explicitly a pro-p-extension with a given Galois group it is convenient to describe the Galois group in term of generators for the quotients of two consecutive terms in a central series.

Let recall how a p-group can be presented by powers and commutators (see [Jo, 13.2]).

Let G be a p-group of order p^n and take a central series

$$G = G_0 \subset G_1 \supset ... \supset G_n = 1$$

such that $|G_i/G_{i-1}| = p$. Take a_i a non trivial element of $G_{i-1} \backslash G_i$ for all $1 \leq i \leq n$. There is a power-commutator presentation of G :

$$G =< a_1, ..., a_n \mid P, C >$$

where the relations are :

$$P \ : \ a_j^p = \prod_{k=j+1}^{n} a_k^{\alpha(j,k)}, \ 1 \leq j \leq n, \ 1 \leq \alpha(j,k) \leq p;$$

$$C \ : \ [a_j, a_i] = \prod_{k=j+1}^{n} a_k^{\beta(1,j,k)}, \ 1 \leq i < j \leq n, \ 1 \leq \beta(1,j,k) \leq p.$$

We call the a_i's the PC-generators (not to confuse with the group minimal set of generators).

The notion of PC-presentation can be extended to pro-p-groups by taking the inverse limit of PC-presented p-groups

Definition 1.1. *Let G be a pro-p-group with the PC-presentation*

$$G = \varprojlim_{k} < a_1, ..., a_k \mid P_k, C_k >,$$

The set $\{a_i\}$ of elements of G is called a set of PC-generators.

C-extensions are defined by the nature of a PC-presentation of their Galois group.

We need to define the lower central series of a group. Let $\gamma_1(G) = G$ and $\gamma_{i+1}(G) = [\gamma_i(G), G]$. This series is strictly decreasing, i.e $\gamma_i(G)/\gamma_{i+1}(G)$ is trivial if and only if $\gamma_i(G)$ is trivial.

Definition 1.2. *The length of an element a in G, is the greater integer i such that*

$$a \in \gamma_i(G).$$

A filtration of a pro-p-group by open subgroups can be defined as follow (for a finitely generated pro-p-group) :

$$P_0(G) = G,\ P_1(G) = G^p[G,G],\ P_{i+1}(G) = P_i^p(G)[P_i(G),G].$$

It is called the lower p-series.

Definition 1.3. *Let G be a pro-p-group with a set of PC-generators*

$$J = \{a_1, a_2 ..., a_k, ...\}.$$

A C-tower in G is a subset noted (a_{n_k}) of J, satisfying

$$a_{n_3} = [a_{n_1}, a_{n_2}],\ \textit{for certain } n_1 \textit{ and } n_2\ ;$$

$$a_{n_4} = [a_{n_3}, a_{n_2}], a_{n_{k+1}} = [a_{n_k}, a_{n_{k-1}}].$$

Definition 1.4. *a. A C-group is an infinite pro-p-group G which has a set of PC-generators I containing a C-tower C, and if N is the subgroup of G PC-generated by $I \setminus C$, then*

$$\bigcap_{g \in G} g^{-1} N g = 1$$

b. A C-extension is a Galois pro-p-extension whose Galois group is a C-group.

Let L/K be a C-extension and G its Galois group. We will note $N(E)/K$ the normal closure of an extension E/K and $M = L^N$ will be called the extension of K defined by the C-tower C. Also we will note K_i the subfields of M/K such that $Gal(K_i/K_{i-1}) =< a_i >$ (where we identify a_i with its image in $Gal(K_i/K_{i-1})$).

We call an extension of C_n-type an extension E/K such that there exists a C-extension L/K such that $L^N = \cup K_i$ as above and $E = K_n$. If E is an extension of C_n-type, we will note $N(E)$ the normal closure of the extension E/K

Remark. The existence of C-extensions will be proved in Section 2.

The purpose of the next lemma is to describe the elements in the Galois group of a C-extension which are not in the C-tower.

Lemma 1.5. *The elements* b_k *in* $Gal(N(K_k)/K_kN(K_{k-1}))$ *are such that*

$$a_k = [b_k, a], \text{ and } b_k \in < [a_{k-1}, a_j], 2 \leq j \leq k_2 >$$

with a *an element in* $Gal(N(K_k)/K)$ *which does not belong to*

$$Gal(N(K_k)/K_kN(K_{k-1})).$$

Proof. The assertion $a_k = [b_k, a]$ comes from the fact that $K_kN(K_{k-1})/K$ is not a normal extension, therefore

$$Gal(N(K_k)/K_kN(K_{k-1}))$$

is a subgroup of $Gal(N(K_k)/K)$ which is not normal. For all b_k in

$$Gal(N(K_k)/K_kN(K_{k-1})),$$

there is an element a in $Gal(N(K_k)/K))$ with a non trivial image in

$$Gal(N(K_{k-1})K_k/K)$$

such that

$$[b_k, a] \notin Gal(N(K_k)/K_kN(K_{k-1}))$$

and as $N(K_k)/N(K_{k-1})$ is a normal extension of K, we have that

$$[b_k, a] = a_k.$$

We recall how a normal subgroup of a p-group is generated (see [Jo]). Let G be a p-group and H be a normal subgroup. Let I be a set of representative in G of elements in G/H (Schreier transversal). Let J be a set of generators of G. A generating set of H (not necessarily minimal) is the following

$$\{tx(\bar{tx})^{-1}, \ x \in J, \ t \in I\}$$

where $\bar{tx}$ is the representative in I of the image of tx in G/H.

The element a_k is a generator of the normal subgroup $H_k = Gal(N_k/N_{k-1})$, but the normal subgroup of H_k not containing a_k define a subextension of N_k/K which is not normal. So there is an element $b_{k,1}$ such that

$$a_k = [b_{k,1}, a] \in Gal(N_k/K)$$

with a a lift of an element in $Gal(N_{k-1}/K)$. If the normal subgroup of H_k not containing $b_{k,1}$ define a subextension of N_k/K which is normal then $\{a_k, b_{k,1}\}$ define a system of generators of the normal subgroup H_k of $Gal(N_k/K)$. If not, there exists an element $b_{k,2}$ such that

$$b = [b, a] \in Gal(N(K)/K)$$

This process is finite because H_k is a finite group. Now we have to show that the set of generators can be taken as in the lemma :

Writing explicitly the commutators we obtain the following equality, by assuming that whenever a commutator was of length greater that the length of a_k it was trivial in $Gal(N_{k-1}K_k/K)$ by the above :

$$a_k = [a_{k-1}, a_{k-2}] = [a_{k-1}, [a_{k-3}, a_{k-4}]]$$
$$= [a_{k-1}^{-1}, a_{k-4}^{-1}][a_{k-4}^{-1}, [a_{k-1}^{-1}, a_{k-4}^{-1}]][a_{k-4}, [a_{k-3}, a_{k-1}]][a_{k-1}^{-1}, a_{k-4}][[a_{k-1}^{-1}, a_{k-4}], a_{k-3}].$$

The elements

$$[a_{k-3}, a_{k-1}] \text{ and } [a_{k-1}, a_{k-4}]$$

are non trivial if and only if a_k is non trivial. Similarly we obtain that $[a_{k-3}, a_{k-1}]$ is non trivial if

$$[a_{k-5}, a_{k-1}] \text{ and } [a_{k-1}, a_{k-4}]$$

are non trivial. And therefore a_k is non trivial if for all $2 \leq i \leq k-2$, $[a_{k-1}, a_i]$ is non trivial. □

Proposition 1.6. *Let L/K be a C-extension with Galois group G. The pro-p-group G is just infinite, i.e. all proper closed normal subgroups of G are open.*

Proof. Let H be a closed normal subgroup of G. If H contains a_i an element in the C-tower, then all a_j, $j > i$ are in H. Indeed, by normality of the subgroup $[a_i, a_{i-1}] = a_{i+1}$ is in H and the result is obtained by induction. In Lemma 1.5, we saw that all the PC-generators of G which are not in the C-tower are elements in the group spanned by $\{[a_{k-1}, a_j], 2 \leq j \leq k_2\}$. If almost all a_i's are is in H, then almost all $[a_{k-1}, a_j]$ is also in H by normality of H and then almost all PC-generator of G is in H. Therefore H is an open subgroup of G.

Now assume that no element of the C-tower is in H. H contains a PC-generator b of G which is not in the C-tower, such that $a_k = [b, a]$ for a certain integer k and a is an element of G. By normality of H a_k is in H and as above H is open.
□

The direct consequence of this result is that it will be rather simple to check if a C-extension is APF, as it is enough to check the unboundedness of the sequence of upper ramification breaks.

Corollary 1.7. *A C-extension L/K is APF if and only if it is deeply ramified, i.e. if the sequence of upper ramification breaks of L/K is unbounded.*

Proof. Let $G = Gal(L/K)$. If L/K is APF then it is deeply ramified. Assume that sequence of upper ramification breaks of L/K is unbounded then every upper ramification group G^u for $u > 0$ is non trivial. By Prop. 1.6 all G^u for $u > 0$ is also open. □

We show that these extensions will give examples of APF extensions where the unboundness of the sequence of upper ramification breaks will not be the consequence of a $\mathbb{Z}_p$-quotient in an open subgroup.

Proposition 1.8. *All open subgroup H of G contains no normal closed subgroup N such that*

$$H/N \simeq \mathbb{Z}_p.$$

Proof. Let H be an open subgroup of G and there exists an element a which is a generator of H and an element of infinite order ([Ze]). But the principle of the proof is to show that for all open subgroup H the derived group $[H, H]$ is of finite index. Therefore there are no quotient of H isomorphic to $\mathbb{Z}_p$ which is an infinite abelian group.

We show that the lower central series are equal to the lower p-series, i.e

$$\forall i,\ \gamma_i(G) = P_i(G) = P_{i-1}(G)^p[P_{i-1}(G), G]$$

We have to show that

$$\gamma_i^p(G) \subset [\gamma_i(G), G].$$

Let a be an element in G of length ℓ. There is a commutator $[b, c]$ of length ℓ such that $a = [b, c]a'$.

$$a^p = [b, c]^p a'^p ...$$

the ... stand for commutators of higher length.

$$[b, c]^p = [b^p, c][b, [b, c]]^{p-1} ...$$

as we have proved in Lemma 1.5 that there are no subgroup of any finite quotient of G generated by an element a_i^p such that the length of a_i^p is the length of a_i. So either $b^p = 1$ or we can decompose again in the same way, and after a finite number of time we obtain that the length of a^p is strictly greater than ℓ.

G being a finitely generated pro-p-group such that G^p is included in the derived group $[G, G]$, this last subgroup is of finite index in G. Let H be an open subgroup of G, there exists an integer i such that

$$\gamma_i(G) \subset H.$$

Therefore

$$[\gamma_i(G), \gamma_i(G)] \subset [H, H].$$

To show that the derived group $[H, H]$ is of finite index in H it is enough to show that for all i, $[\gamma_i(G), \gamma_i(G)]$ is of finite index in G.

There is an inclusion

$$[\gamma_i(G), \gamma_i(G)] \subset \gamma_{2i}(G).$$

The quotient

$$\gamma_{2i}(G)/[\gamma_i(G), \gamma_i(G)]$$

contains only elements in

$$[G/\gamma_2(G), \gamma_{2i-1}(G)/\gamma_{2i}(G)], [\gamma_2(G)/\gamma_3(G), \gamma_{2i-2}(G)/\gamma_{2i-1}(G)], ...,$$

All these groups being finite, the quotient

$$\gamma_{2i}(G)/[\gamma_i(G), \gamma_i(G)]$$

is finite. The subgroup $\gamma_{2i}(G)$ is of finite index in G, and so is $[\gamma_i(G), \gamma_i(G)]$. For every open subgroup H of G, the derived group $[H, H]$ contains a subgroup which is open in G and we have proved the result. □

The last result of this section shows that if a C-tower is APF, this is not due to the result of Sen on p-adic Lie groups.

Proposition 1.9. *The Galois group of a C-extension is not a p-adic analytic group.*

Proof. It is enough to show that there are open subgroups with an arbitrary large number of generators, i.e we show that the pro-p-group G is not of finite rank (see [DDMS] Prop. 3.1 and Cor. 8.34). Consider the subgroup H of G containing

$$a_2, a_3, ..., a_{2k-1}, a_{2k+1}, a_{2k+2}, ...$$

ie. for $i < 2k + 1$ even, a_i is not in the subgroup and for any $i \geq 2k + 1$ a_i is in the subgroup. Then the elements

$$a_2, a_3, ..., a_{2k-1}, a_{2k+1}$$

are topological generators of the open subgroup H. Indeed, none of this elements are in the Frattini subgroup $\phi(H)$ or are equal to another one modulo an element in $\phi(H)$:

$$a_{2i+1} = [a_{2i}, a_{2i-1}], a_{2i} \notin H, a_{2i-1} \in H.$$

□

2. There exist APF C-extensions

In this section we assume that K is a finite extension of $\mathbb{Q}_p$ of degree at least 3. We construct a totally ramified C-extension and show that there exist C-extensions of K which are APF.

Proposition 2.1. *Let K be a finite extension of $\mathbb{Q}_p$ of index of ramification $e(K)$. For all $n > 0$ there exists an extension of C_n-type with maximal upper ramification break satisfying*

$$u_n > ne(K) - v_n \text{ where } \lim_{n\to\infty} v_n < \infty.$$

First we prove a lemma.

Lemma 2.2. *i. Let K_1/K and E/K_1 be cyclic extensions of order p such that E is not normal over K. Then there exists a cyclic extension K_2/K of order p such that K_2E/K is a normal extension of order p^3 with Galois group having the presentation :*

$$Gal(K_2E/K) =< a_1, a_2 \mid a_1^p = a_2^p = [a_1, a_2]^p = 1, [a_1, [a_1, a_2]] = [a_2, [a_1, a_2]] = 1 >$$

ii. Let K_1/K be a cyclic extension of order p and E/K_1 an extension of C_n-type. We have one of the following cases :

a. there exist a cyclic extension K_2/K of order p such that K_2E/K is a normal extension of C_{n+2}-type ;

b. there exists two cyclic extensions K_2/K and K_3/K of order p such that $K_2K_3N(E)/K$ is a normal extension and if we note b and c a lift of the generators $Gal(K_2/K)$ and $Gal(K_3/K)$ in

$$Gal(K_2K_3N(E)/K)$$

then we have the relations in $Gal(K_2K_3N(E)/K)$

$$[a, b] = a_1\alpha_1 \quad ; \quad [a, c] = a_2\alpha_2$$

where $(a_1\alpha_1, a_2\alpha_2)$ define a minimal system of generators of

$$Gal(K_2K_3N(E)/K_1K_2K_3)$$

and a is a lift of a generator of $Gal(K_1/K)$.

Proof. i. Let take $E(K)$ the normal closure of E/K and consider a generator b of $Gal(E(K)/K)$ which has a trivial image in $Gal(K_1/K)$. We note a (resp. c) a lift of a generator of $Gal(K_1/K)$ (resp. $Gal(E/K_1)$) in $Gal(E(K)/K)$. Then the subgroup generated by b is not a normal subgroup. It means that one can choose the generators such that there is one of the relations which is

$$[a, b] = c \text{ or } [b, c] = a.$$

We note that the subgroup generated by b and c is normal so the second relation is not possible and we get the result.

ii. We obtain two extensions K_2/K and K_3/K by considering two cyclic independent subextensions of E/K_1 as in case i. Let call b and c the two generators obtained. There are two possibilities : either K_2 and K_3 are linearly independent in which case we are in case b. or $c = b^i$ $i \neq 1$ and we are in case a, indeed we have that $a_1 = [a, b]$ and

$$a_2 = [a, b^i] = [a, b]^i[[a, b], b]$$

and therefore

$$[a_2, a_1] = [[[a, b], b], [a, b]].$$

□

Proof of Proposition 2.1. Let show first that for all K non trivial extension of $\mathbb{Q}_p$, there exists an extension E_3/K of C_3-type satisfying the condition on the ramification. Let take an extension K_1/K with ramification break $i_1(3)$ and an extension E/K_1 with ramification break i all of them cyclic of order p such that $i > i_1(3)$ and we put

$$u_3 = i_1(3)\frac{p-1}{p} + \frac{i}{p}.$$

Moreover we want E not to be normal over K. For this it enough to take i such that p does not divide $i - i_1$. By Lemma 2.2, there exists a normal extension E_3/K of C_3 type for which the bigger upper ramification break is u_3.

Let K_1/K be a cyclic extension of order p with ramification breaks $i_1(5)$. We showed that there exists an extension E_3/K_1 with bigger upper ramification break u_3. We put

$$u_5 = i_1(5)\frac{p-1}{p} + \frac{u_3}{p}.$$

By Lemma 2.2 we have two possibilities either $c = b^i$ and we have the result or $Gal(E(K)/K)$ is a p-group with exactly three generators.

We show for all n that $Gal(E(K)/K)$ contains a subgroup N such that the subextension $E(K)^N/K)$ is the subextension of an extension of C_n-type.

Take the normal subgroup N of $Gal(E(K)/K)$ generated by $c[a,b]$. The group N also contains

$$a^{-1}c[a,b]a = c[c,a][a,b][[a,b],a] \text{ and}$$

$$b^{-1}c[a,b]b = c[c,b][a,b][[a,b],b]$$

which will define relations in $Gal(E(K)^N/K)$.

The map

$$\begin{array}{ccc}
Gal(E(K)/K) & \to & Gal(E(K)/K)/N \\
a & \to & \bar{a} \\
b & \to & \bar{b} \\
c & \to & [\bar{b},\bar{a}] \\
a_1 & \to & [\bar{a},\bar{b}] \\
a_2 & \to & [\bar{a},[\bar{a},\bar{b}]] \\
b^{-1}c[a,b]b & \to & 1 \\
a^{-1}c[a,b]a & \to & 1 = [\bar{b},\bar{a}][[\bar{b},\bar{a}],\bar{a}][\bar{a},\bar{b}][[\bar{a},\bar{b}],\bar{a}] \\
\ldots & &
\end{array}$$

shows that $E(K)^N/K$ is the subextension of an extension of C_n-type because the relation

$$[\bar{b},\bar{a}][[\bar{b},\bar{a}],\bar{a}][\bar{a},\bar{b}][[\bar{a},\bar{b}],\bar{a}] = 1$$

is trivial :

$$a_3^{-1}[a_3^{-1},a_2]a_3a_4 = a_2^{-1}a_3^{-1}a_2a_3a_4 = a_2^{-1}a_3^{-1}a_3a_2a_4^{-1}a_4 = 1$$

Let consider the extension of C_n-type which has $E(K)^N$ as a subextension. We proceed with the construction. The bigger ramification break in that extension is u defined above

Now we proved the result for $n = 5$ and we assume we can show it for any odd integer n and show that then for all finite extension K of $\mathbb{Q}_p$ there exists an extension of C_{n+2}-type with bigger upper ramification break u_{n+2}.

Let K_1/K be a cyclic extension of order p with ramification break $i_1(n+2)$. By assumption there exists an extension of K_1 of C_n-type with biggest upper ramification break u_n then put

$$u_{n+2} = i_1(n+2)\frac{p-1}{p} + \frac{u_n}{p}.$$

By Lemma 2.2, the situation is similar to the case $n = 5$. Either there exists an extension K_2/K such that E_nK_2/K is of C_{n+2} type or we have to take a quotient of $Gal(E_n(K)/K)$ by the same normal subgroup as in the previous case.

It remains to show that the ramification breaks $i_1(3), i_1(5), ..., i_1(2n+1), ...$ can be taken such that upper ramification breaks $u_3, .., u_n, ...$ satisfy the condition in the proposition, i.e there exists a convergent sequence v_n such that

$$u_n > ne(K) - v_n.$$

For $n = 3$ take any $i_1(3)$ and i.

Then for $n = 5$, we take

$$i_1(5) = \frac{pe(K)}{p-1} - \varepsilon_5$$

where ε_5 is an integer, such that

$$u_5 = e - \frac{p-1}{p}\varepsilon_5 + \frac{u_3}{p}.$$

For $n = 7$, we take

$$i_1(7) = \frac{pe(K)}{p-1} - \varepsilon_7$$

such that

$$u_7 = e - \frac{p-1}{p}\varepsilon_7 + \frac{u_5}{p} = e - \frac{p-1}{p}\varepsilon_7 + \frac{pe(K)}{p} - \frac{p-1}{p^2}\varepsilon_5 + \frac{u_3}{p^2}$$

$$= 2e - [\frac{p-1}{p}(\varepsilon_7 + \frac{\varepsilon_5}{p}) - \frac{u_3}{p^2}].$$

Note that for $n+1$ we have that the value $e(K_1) = pe(K)$

By taking for all odd integer k,

$$i_k = \frac{pe(K)}{p-1} - \varepsilon_k$$

we obtain

$$u_{2n+1} = (n-1)e - [\frac{p-1}{p}(\varepsilon_{2n+1} + \frac{\varepsilon_{2n-1}}{p} + ... + \frac{\varepsilon_5}{p^{n-1}}) - \frac{u_3}{p^{n-1}}].$$

Take all the ε_k in a bounded interval, then

$$v_n = [\frac{p-1}{p}(\varepsilon_{2n+1} + \frac{\varepsilon_{2n-1}}{p} + ... + \frac{\varepsilon_5}{p^{n-1}}) - \frac{u_3}{p^{n-1}}].$$

is a convergent sequence.

□

Corollary 2.3. *There exist APF C-extensions.*

Proof. Let L/K be a C-extension and note $\{a_i\}$ a C-tower defining its Galois group. We assume that this extension is not APF. Take a family E_n of extensions of K linearly independent of L/K such that for all n E_n/K is an extension of C_n-type and its greater upper ramification break $\tilde{u}_n$ satisfies

$$\tilde{u}_n \geq ke \text{ where } n = 2k \text{ or } n = 2k - 1.$$

This family exists by Proposition 2.1 and one can check that the extensions taken are linearly independent of L/K by checking that the compositum of the L^{ab}/K and the subextensions $E_{n,1}/K$ of E_n of order p^2 which is not normal remains not normal.

We will note $\{\tilde{a}_{i,n}\}$ the C-tower defining the extension E_n/K.

Let u_n be the sequence of upper ramification breaks in L/K. This sequence is convergent as we assumed that the extension is not APF. We modify the extension L/K in order to obtain a C-extension $\tilde{L}/K$ which is APF. Consider the compositum M of L with all the finite extensions $N(E_n)/K$ and take $\tilde{L}$ the fixed field of M by the normal subgroup generated by $a_1\tilde{a}_{1,n}^{-1}$ and $a_2\tilde{a}_{2,n}^{-1}$ for all n. We show that this is a well defined C-extension and that it is APF, i.e. that the sequence of upper ramification breaks in $\tilde{L}/K$ is unbounded (see Cor. 1.7).

The Galois group $Gal(\tilde{L}/K)$ is defined by the surjective map

$$\begin{array}{ccccc}
Gal(M/K) & \rightarrow & Gal(\tilde{L}/K) & \\
a_1 & \rightarrow & \bar{a}_1 & \\
a_2 & \rightarrow & \bar{a}_2 & \\
a_{1,n} & \rightarrow & \bar{a}_1 & \forall n > 2 \\
a_{2,n} & \rightarrow & \bar{a}_2 & \forall n > 2 \\
a_k & \rightarrow & \bar{a}_k(\bar{a}_1, \bar{a}_2) & \forall n > 2 \\
a_{k,n} & \rightarrow & \bar{a}_k(\bar{a}_1, \bar{a}_2) & \forall n > 2
\end{array}$$

where $\bar{a}_k(\bar{a}_1, \bar{a}_2)$ is the k^{th} element is the C-tower generated by $\bar{a}_1$ end $\bar{a}_2$. This a group homomorphism. So $\tilde{L}/K$ is a C-extension.

The upper ramification breaks $\bar{u}_k$ in $\tilde{L}/K$, for k large enough satisfy:

$$\bar{u}_k \geq Sup\{u_{L/K}(a_k), u_{E_n/K}(a_{k,n}), \forall n \geq k\}.$$

Indeed, this is not the case if for all $n \geq k$

$$u_{L/K}(a_k) = u_{E_n/K}(a_{k,n}) \text{ and } u_{M/K}(a_k a_{k,n}) > u_{L/K}(a_k).$$

If L/K is not APF, for k big enough

as the first sequence is convergent and

$$u_{E_k/K}(a_{k,k}) > ([\frac{k-1}{2}]-1)e(K).$$

With the hypothesis made on the greater upper ramification break in the extension E_n/K we obtain that for n big enough

$$\bar{u}_n \geq ([\frac{n-1}{2}]-1)e(K).$$

and therefore the extension $\tilde{L}/K$ is APF. $\square$

3. THERE EXIST NON ARITHMETICALLY PROFINITE C-EXTENSIONS

We want to show that unlike extensions whose Galois group is a p-adic Lie group, the structure of the Galois group of C-extensions does not implies that the extensions is APF. Indeed, we show how to construct a C-extension which is not APF.

Proposition 3.1. *There exist C-extensions which are not deeply ramified .*

We construct a C-extension L/K such that L^N/K has bounded ramification.

Let L/K be a C-extension such that $L^N = \cup K_n$ as defined in Section 1. Consider the (not normal) closed subgroup H of $G = Gal(L/K)$ such that only the elements of the C-tower $a_{2i+1}, i \geq 0$ are not in H and take $M' = L^H$ the fixed field of L by H. Then M'/K is an infinite extension with $M' = \cup M_n$ such that for all n M_n/M_{n-1} is a cyclic extension of order p and M_n is not a normal extension of M_{n-2}. Moreover we assume that for all n we have that the ramification breaks satisfy :

$$p \nmid (i(M_n/M_{n-1}) - i(M_{n-1}/M_{n-2})).$$

This is possible to obtain by the methods used in Section 2.

We are going to construct a family of extensions E_n/K of C_n-type for all $n > 1$ such that there exists a subextension of $L \cup (\cup N(E_n))$ which is a non APF C-extension.

First we show that we can construct a tower of extensions E'_i/K such that for all i the extension

$$M_iE'_{i-1}E'_iM_{i-1}/M_{i-1}E'_{i-1}$$

contains a cyclic subextension of order p with a given ramification break

$$i_n < i(M_iE'_{i-1}/M_{i-1}E'_{i-1}).$$

We show a general lemma on the existence of Galois extension of order p^2 with given ramification breaks :

Lemma 3.2. *Let L/K be a cyclic extension of order p with ramification break i. Let j and s be two positive integers not divisible by p such that if $s \leq i$ then $s \leq j$ and $i + p^{-1}(s-i) \leq j$ otherwise ; and $j \leq \dfrac{pe}{p-1}$ where e is the absolute index of ramification of K. If $i \neq j$, take $s = h_{L/K}(j)$. There exists a cyclic extension L'/K of order p with ramification break j such that the ramification break of the cyclic extension LL'/L is s.*

Proof. Suppose that $i \neq j$.

There exists a cyclic extension K_1/K of order p totally ramified with ramification break j if and only if $j \leq \dfrac{ep}{p-1}$ ([FV, Prop. III2.5]). As $i \neq j$, the ramification in LL'/L is $s = h_{L/K}(j)$.

Suppose that $i = j$. Let λ be a root of the polynomial $X^p - X - \alpha$ with $v_K(\alpha) = -i$ and $L = K(\lambda)$ and $<\tau>= Gal(L/K)$. We show that there exists an element β in K, $v_K(\beta) = -i$ and μ the root of the polynomial $X^p - X - \beta$ such that there is an element A in $K(\lambda, \mu)$ stable by $\tau\sigma$ where σ generates $Gal(K(\mu)/K)$ satisfying the conditions :

. $v_{K(\lambda,\mu)^{\tau\sigma}}(A) = -s$

. $v_{K(\lambda,\mu)^{\tau\sigma}}(\sigma(A) - A - 1) > 0$.

It is enough to show that $i(Gal(K(\lambda,\mu)^{\tau\sigma}/K)) = -s$ and therefore

$$i(Gal(K(\lambda,\mu)/L)) = -s.$$

If $p = 2$ take

$$A = \mu\tau(\lambda) + \lambda\sigma(\mu),$$

if $p > 2$ take

$$A = \sum_{0 \leq i < j \leq p-2} \tau^i(\lambda)\sigma^j(\mu) - \tau^j(\lambda)\sigma^i(\mu).$$

First we show that $\tau\sigma(A) = A$.

$$\tau\sigma(A) = \sum_{0 \leq i < j \leq p-2} \tau^{i+1}(\lambda)\sigma^{j+1}(\mu) - \tau^{j+1}(\lambda)\sigma^{i+1}(\mu).$$

We conclude by the fact that $Tr_{K(\mu)/K}(\mu) = Tr_{K(\lambda)/K}(\lambda) = 0$ which allows to replace $\tau^{p-2+1}(\lambda)$ by $-(\sum_{i=0}^{p-2} \tau^i(\lambda))$ and $\sigma^{p-2+1}(\mu)$ by $-(\sum_{i=0}^{p-2} \sigma^i(\tau))$.

For $p = 2$, the result is clear.

We show that the second condition is always valid. We have that

$$v_{K(\lambda)}(\tau^j(\lambda) - \tau^i(\lambda) - 1) > 0$$

and

$$v(\sigma^j(\mu) - \sigma^i(\mu) - 1) > 0$$

for $j > i$. For $p > 2$

$$\sigma(A) - A = \sum_{0 \leq i < j \leq p-2} \tau^i(\lambda)(\sigma^{j+1}(\mu) - \sigma^j(\mu)) - \tau^j(\lambda)(\sigma^{i+1}(\mu) - \sigma^i(\mu))$$

$$\sigma(A) - A = \sum_{0 \leq i < j \leq p-2} (\tau^i(\lambda) - \tau^j(\lambda) + ...)$$

where the dots mean terms of higher valuation. The above sum is then equal to

$$\sigma(A) - A = 1 - p + B$$

where B is of positive valuation, which gives the result.

In the case $p = 2$ one has

$$\sigma(A) - A = (\sigma(\mu) - \mu)(\tau(\lambda) - \lambda),$$

so we can conclude in a similar way.

Let $M = K(\lambda, \tau)^{\tau\sigma}$. By a similar method as above we obtain :

$$v_M(A) = v_M(\sum_{0 \leq i < j \leq p-2} \tau^i(\lambda)(\sigma^j(\mu) - (\sigma^i(\mu)) - \sigma^i(\mu)(\tau^j(\lambda) - \tau^i(\lambda)))$$

$$= v_M(\sum_{0 \leq i < j \leq p-2} (\tau^i(\lambda) - \sigma^i(\mu))).$$

From the equality of the valuations

$$v_M(\tau^i(\lambda) - \sigma^i(\mu)) = v_M(\tau^i(\lambda)^p - \sigma^i(\mu)^p - \alpha + \beta)$$

we deduce that $v_M(\tau^i(\lambda) - \sigma^i(\mu)) = -s$ if

$$(*) \qquad v_K(\alpha - \beta) = -s$$

and if this last equality is valid then $v_M(A) = v_K((p-2)(\alpha - \beta)) = -s$. So take $\beta = \alpha + \pi_K^{-s}$ and we have proved the existence of the extension $K(\mu)$ with the required properties.

For $p = 2$ we obtain the condition $v_M(A) = -s$ if $v_K(\alpha + \beta) = -s$. □

The lemma gives the existence of an extension $F/M_{i-1}E'_{i-1}$ satisfying the property, but we have to make sure that this correspond to an extension of E'_{i-1}. For this take an element β in E'_{i-1} such that (see (*) in the proof of the lemma)

$$v_{M_{i-1}E'_{i-1}}(\alpha - \beta) = -s.$$

It remains to check that the extension obtained is not normal over E'_{i-1}. This is clear with the assumption on the lower ramification breaks of the extensions M_i/M_{i-1} made at the beginning of this section.

We have a tower of extensions E'_i such that E'_{i+1}/E'_{i-1} is not normal. We construct the extensions E_n/K in a similar way that we constructed the extensions E_n/K in the previous section.

The extension E_3/K is obtained as in Section 3 by taking the normal closure of E'_2/K. Then we obtain E_5, we start by taking the normal closure of E'_3/E'_1 to obtain a Galois extension F_5/E'_1 which is not normal over K and then take the the normal closure of this extension. We get E_5/K as in Section 2.

We can repeat this process for all n and obtain the required family of extensions. Now we look at the ramification in the subextension L' of $L \cup (\cup N(E_n))$ which is the fixed field of $L \cup (\cup N(E_n))$ by the normal subgroup generated by $a_1\tilde{a}_{1,n}^{-1}$ and $a_2\tilde{a}_{2,n}^{-1}$ for all n keeping the notation of Section 2. We saw that it was a C-extension. We note that for all i, n and m, $\tilde{a}_{2i,n} = \tilde{a}_{2i,m}$. In the construction of the tower of E'_n, we can choose the ramification breaks i_n to be small enough so that the sequence of upper ramification breaks in the extension L'/K converges.

4. Construction of some APF extensions whose Galois group is not a p-adic Lie group

First we discuss deeply ramified extensions. We know that all totally ramified extensions whose Galois group contains an open subgroup with a $\mathbb{Z}_p$-quotient are deeply ramified. In this paper we have shown that :

Proposition 4.1. *i. All pro-p-groups that contains a C-tower can be realized as the Galois group of a deeply ramified extension.*

ii. The fact that the Galois group of a totally ramified extension contains a C-tower is not enough to say that the extension is deeply ramified.

Now we look at arithmetically profinite extensions.

The result obtained in Section 2 can be extended to construct more APF Galois pro-p-extensions

Proposition 4.2. *Let E/K be a totally ramified Galois pro-p-extension such that*

. $Gal(E/K)$ is a pro-p-group over two generators ;

. E is the compositum of a finite number of extensions E_i/K each of them containing a subextension L_i/K such that E_i/L_i is a C-extension.

Then for all subextension L/K of E/K which is infinite, $Gal(N(L)/K)$ contains a subgroup which is generated by a C-tower (i.e there is a finite extension K'/K such that L/K' is a C-extension.

Proof. Let call

$$\{a_{k,i}, i > 0\} \text{ for } 1 \le k \le n$$

the C-towers generating a lift of $Gal(E_k/L_k)$ in $G = Gal(E/K)$

Let H be a normal subgroup of G of infinite index. We show that at least one C-tower is not contained in H. Suppose that

$$a_{k,i} \in H$$

for a certain pair (k, i) and

$$a_{k,i-1} \notin H.$$

then as H is normal we have

$$a_{k,i-1}^{-1} a_{k,i} a_{k,i-1} = a_{k,i} a_{k,i+1} \in H$$

and therefore

$$a_{k,i+1} \in H.$$

So as there are only a finite number of C-towers and nothing else there is a C-tower which has a non trivial image in G/H.

We show that there is an open normal subgroup of G/H which is generated by a C-tower. Let $\{\bar{a}_i\}$ be the image of a C-tower in G/H. As G is the union of a finite number of C-towers, there are a finite number of $\bar{a}_i$ such that there exists an element $\bar{b}$ in G/H which is not in the subgroup generated by $\{\bar{a}_i\}$ which does not commute with $\bar{a}_i$. So there exists i_0 such that the subgroup generated by the C-tower $\{\bar{a}_i,\ i > i_0\}$ is an open normal subgroup of G/H.

□

The sequence of upper ramification breaks in a quotient depends on the relations that define this quotient :

Proposition 4.3. *Let E/K be as in Prop. 4.2. Let H be a closed normal subgroup of infinite index. Let $\{\bar{a}_i\}$ be a C-tower in G/H. The upper ramification breaks of the elements in $\{\bar{a}_i\}$ for i big enough are given by :*

$$u_{G/H}(\bar{a}_i) = Max\{u_G(a_{k,i}), \varphi(a_{k,i}) = \bar{a}_i\}.$$

Proof. To see this we have to show that if there exists a C-tower $\{a_{k,i}\}$ which is send to $\{\bar{a}_i\}$ in G/H, then either no other C-tower is mapped to this one or if there is then $a_{l,i}...$ is mapped to $\bar{a}_i$ where the dots stand for elements $a_{l,j}$, $j > i$.

This comes from the fact that if

$$a_{k,i} \prod_{l \in I, j \in J} a_{l,j} \in H$$

then the towers $\{a_{l,i}\}$ have a non trivial image in G/H if an elements that does not commutes with $a_{l,j}$ is mapped to $\bar{a}_i$.

We can show that this element can only be $a_{l,j-1}....$ And by going down until there are no more elements in G/H which commute with one of the elements, we see that we must have $i = j$

□

We have the following result on the ramification in the extension L/K :

Corollary 4.4. *If for all C-towers in G we have that*

$$u_G(a_{k,i}a_{l,i}) = Inf\{u_G(a_{k,i}), u_G(a_{l,i})\}$$

and all the C-extensions E_i/L_i are APF. Then E/K is APF.

A consequence of this result is that we can construct an APF extension with a Galois group as in Proposition 4.3

Proposition 4.5. *There exists extension E/K as in Proposition 4.2 which are arithmetically profinite.*

Proof. To construct extensions E/K which are APF we use the same method as in Section 2. By Corollary 4.4, it is enough to take all the C-towers having distinct upper ramification breaks.

Take an extension E/K as in Prop. 4.2. We can modify it so that all the C-extensions E_i/L_i are APF as done in Section 2. In order to have the condition of Cor. 4.4 we have to modify them so that they have all distinct upper ramification breaks sequences which can be done because there is a finite number of C-extensions to consider. □

A similar result can be obtained for the compositum of infinitely many C-extensions over the same two generators if we restrict to such group in which every elements in a C-tower commutes with almost all elements in G. But even in that case there are infinite Galois subextensions whose Galois group does not contain any C-tower. Namely one can take the normal subgroup that contains all the $a_{k,i}$ for i greater that a given i_0, for all k. We can show that there no other Galois subextensions than those and the one which we looked at the beginning of that section but we have to make sure that such subextensions are also APF.

We discuss briefly how to generalize the construction made in Section 2, in order to obtain APF extensions of this type.

Let K_1/K be a cyclic extension of order p and let $K_2/K_1, ..., K_p/K_1$ be $p-1$ distinct cyclic extensions of K_1 with different ramification breaks such none of them are normal over K. Then the result obtained in Lemma 2.1 extend and therefore there is a quotient of the normal closure of the extension $K_2K_3...K_p/K$ whose Galois group is PC-generated by $a_1, a_2, a_3, ...a_{p+1}$ with $a_3 = [a_2, a_1]$ and $a_i = [a_{i-1}, a_1]$ for $i > 3$.

Now we have to check that, in this type of construction, we can take the ramification to be hight enough so that the whole extension is APF. By considering the number of generators in an open subgroup of a free pro-p-group with two generators, the bound on the upper ramification and the fact that we want the upper ramification breaks to be unbounded, we can't simplify our problem by taking all ramification distinct because there would not be enough integer available. But as seen in Section 3, we can control the extensions so that the ramification breaks would not collapse in certain quotients.

References

[CG] J. Coates & R. Greenberg Kummer theory for abelian varieties over local fields, Invent. Math. **124** (1996), 129–174.

[DDMS] J.D. Dixon, M.P.F. du Sautoy, A. Mann, D. Segal, *Analytic pro-p-groups,*, Cambridge studies in advanced mathematics, 2nd ed, Cambridge Univ. Press **61** (1999).

[DuFe] M.P.F. du Sautoy & I.B. Fesenko, *Where the wild things are : ramification groups and the Nottingham group*, New horizons in pro-p-groups,du Sautoy and A. Shalev (eds.) , Progress in Math. **184** (200), 287–328.

[Fe] I.B. Fesenko, *On just infinite pro-p-groups and arithmetically profinite extensions of local field*, J. reine angew. Math. **517** (1999), 61–80.

[FV] I. Fesenko & S. Vostokov, *Local Fields and their Extensions, a Constructive Approach*, Transl. of Mathematical Monographs **121**.

[FW] J.-M. Fontaine & J.-P. Wintenberger, *Corps de normes de certaines extensions algebriques de corps locaux*, Comptes rendus **288 série A** (1979), 441–444.

[Jo] D.L. Johnson, *Presentation of groups*, LMS Student Texts **15** (1990).

[Sen] S. Sen, *Ramification in p-adic Lie extensions*, Invent. Math. **90** (1972), 44–50.

[Ser] J.-P. Serre, *Corps Locaux*, Hermann (1968).

[Ze] E.I. Zelmanov, *On periodic compact groups*, Israel J. Math. **77** (1992), 83–95.

UNIVERSITY OF CAMBRIDGE, DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, CENTRE FOR MATHEMATICAL SCIENCES, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB, UK

E-mail address: `O.Sauzet@dpmms.cam.ac.uk`